

PCR protocol

RBRC 03390

Strain name ApoE3-3 KO

Primer	primer_name	primer_seq	length	
1	oIMR0180	GCCTAGCCGAGGGAGAGCCG	20	JAX Genotyping Protocol
2		CACCAGAGGGCCAGAGAGGTGC	22	RIKEN BRC
3	Neo-P1	GCCTGCTTGCCGAATATCATGGTGGAAAAT	30	Depositor's primer


Reaction Components

Enzyme TOYOBO KOD FX (Code.No : KFX-101)

	Conc	Run 1	Run 2	Run 3	Run 4
H2O		1.8			
PCR buffer	2 X	10.0			
dNTP	2.0 mM	4.0			
primer 1	10 pmol/ μ l	1.0			
primer 2	10 pmol/ μ l	1.0			
primer 3	10 pmol/ μ l	1.0			
DNA		1.0			
Taq	1 U/ μ l	0.2			
total		20			

Reaction Conditions

step	temp (°C)	time (sec)	note
1	94	120	
2	98	10*	
3	63	30*	
4	68	90*	*Repeat for 30 cycles



Marker
1 kb DNA ladder (Invitrogen, Cat. 15615-024)

Gel
1% agarose in 1 X TAE (100 V, 30 min)

PCR products				
	primer_set	product_size	product	
	primer 1	primer 2	1,344 bp	WT
	primer 2	primer 3	ca 800 bp	KI

Comment