

RBRC 06574

Strain name B6.Cg-Tg(Chx10-cre/ERT2)G7Tfur

| Primer | primer_name | primer_seq | length | |
|--------|-------------|---------------------------|--------|-------------------------|
| 1 | | GTCTCCTAGCCTTTGCGTTCAGAC | 24 | RIKEN BRC |
| 2 | Cre3P | TTCGGCTATACGTAACAGGG | 20 | |
| 3 | oIMR0042 | CTAGGCCACAGAATTGAAAGATCT | 24 | JAX Genotyping Protocol |
| 4 | oIMR0043 | GTAGGTGGAAATTCTAGCATCATCC | 25 | JAX Genotyping Protocol |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

Reaction Components


Enzyme TaKaRa Ex Taq (Cat. No. RR001A)

| | Conc | Run 1 | Run 2 | Run 3 | Run 4 |
|--------------|------------------|-----------|-------|-------|-------|
| H2O | | 11.3 | | | |
| PCR buffer | 10 X | 2.0 | | | |
| dNTP | 2.5 mM | 1.6 | | | |
| primer 1 | 10 pmol/ μ l | 1.0 | | | |
| primer 2 | 10 pmol/ μ l | 1.0 | | | |
| primer 3 | 10 pmol/ μ l | 1.0 | | | |
| primer 4 | 10 pmol/ μ l | 1.0 | | | |
| DNA | | 1.0 | | | |
| Taq | 5U/ μ l | 0.1 | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| total | | 20 | | | |

Reaction Conditions

| step | temp (°C) | time (sec) | note |
|------|-----------|------------|-----------------------|
| 1 | 94 | 180 | |
| 2 | 94 | 30* | |
| 3 | 60 | 30* | |
| 4 | 72 | 60* | *Repeat for 30 cycles |
| 5 | 72 | 300 | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

Standard PCR



Marker

1 kb DNA ladder (Invitrogen, Cat. 15615-024)

Gel

1% agarose in 1 X TAE (100 V, 30 min)

PCR products

| | primer_set | product_size | product |
|----------|------------|--------------|-------------------------|
| primer 1 | primer 2 | ca. 1000 bp | Tg |
| primer 3 | primer 4 | 324 bp | Internal control (IL-2) |
| | | | |
| | | | |

Comment